Naval Research Laboratory

Washington, DC 20375-5000 NRL Report 8967 July 28, 1986

UNCLASSIFIED

An Incomplete Lipschitz-Hankel Integral of K_0

Part I

ALLEN R. MILLER

Engineering Services Division

Approved for public release; distribution unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED	1b RESTRICTIVE MARKINGS
2a SECURITY CLASSIFICATION AUTHORITY	3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release, distribution unlimited.
2b DECLASSIFICATION/DOWNGRADING SCHEDULE	
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Report 8967	5 MONITORING ORGANIZATION REPORT NUMBER(S)
6a NAME OF PERFORMING ORGANIZATION Naval Research Laboratory — 6b. OFFICE SYMBOL (If applicable) 2303.1	7a NAME OF MONITORING ORGANIZATION
6c. ADDRESS (City, State, and ZIP Code) Washington, D.C. 20375-5000	7b. ADDRESS (City, State, and ZIP Code)
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Research Laboratory — 8b. OFFICE SYMBOL (If applicable) 2303.1	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER
8c. ADDRESS (City, State, and ZIP Code) Washington, D.C. 20375-5000	10. SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. / PROJECT NO. / TASK NO / WORK UNIT ACCESSION NO.

11 TITLE (Include Security Classification)
An Incomplete Lipschitz-Hankel Integral of K_0, Part I

12 PERSONAL AUTHOR(S)
Miller, Allen R.

13a TYPE OF REPORT	13b. TIME COVERED	14. DATE OF REPORT (Year, Month, Day)	15. PAGE COUNT
Interim	FROM 1/85 TO 1/86	1986 July 28	15

16. SUPPLEMENTARY NOTATION

17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)
FIELD	GROUP	SUB-GROUP	Incomplete Lipschitz-Hankel integrals Kampé de Fériet functions Special functions

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

An incomplete Lipschitz-Hankel Integral of K_0 and related integrals are given in terms of elementary, cylindrical, and Kampé de Fériet functions. Some of the properties of these Kampé de Fériet functions are derived.

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS	21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED
22a NAME OF RESPONSIBLE INDIVIDUAL Allen R. Miller	22b. TELEPHONE (Include Area Code) (202)767-2215 — 22c. OFFICE SYMBOL Code 2303.1

DD FORM 1473, 84 MAR — 83 APR edition may be used until exhausted. All other editions are obsolete — SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

CONTENTS

INTRODUCTION 1

PRELIMINARY DEFINITIONS 1

SOME ELEMENTARY PROPERTIES OF $M[\alpha, \beta; \gamma, \delta; x, y]$ 2

SOME ELEMENTARY PROPERTIES OF $L[\alpha, \beta; \gamma, \delta; x, y]$ 5

A CLOSED FORM FOR $K_{e_0}(a, z)$ 6

KING'S INTEGRAL 9

A DISTRIBUTION FOR THE ELEVATION OF A SINE WAVE 11

SOME INTEGRALS RELATED TO $K_{e_0}(a, z)$ 12

CONCLUSIONS 13

REFERENCES 13

UNCLASSIFIED

AN INCOMPLETE LIPSCHITZ-HANKEL INTEGRAL OF K_0
PART I

INTRODUCTION

An incomplete Lipschitz-Hankel integral of cylindrical functions of order zero, C_0, may be defined by

$$C_{e_0}(a, z) \equiv \int_0^z e^{at} C_0(t)\, dt$$

Of interest in applications are the functions $J_{e_0}(a, z)$, $I_{e_0}(a, z)$, and $N_{e_0}(a, z)$ where J denotes the Bessel function of the first kind, I denotes the modified Bessel function, and N denotes the Bessel function of the second kind or Neumann function. $J_{e_0}(a, z)$ and $N_{e_0}(a, z)$ occur in problems in the theory of diffraction in optical apparatus [1, p. 227]. The function $I_{e_0}(a, z)$ plays an important role in the study of oscillating wings in supersonic flow and arises in the study of resonant absorption in media with finite dimensions [1, p. 195].

In this report we are interested in

$$K_{e_0}(a, z) \equiv \int_0^z e^{at} K_0(t)\, dt \tag{1}$$

where K denotes the MacDonald function or Bessel function of imaginary argument. We shall show that $K_{e_0}(a, z)$ can be written in closed form in terms of elementary functions, K_0, K_1, and Kampé de Fériet double hypergeometric functions. As an application it shall be shown that $K_{e_0}(a, z)$ occurs when the statistical distribution of the maxima of a random function is applied to the amplitude of a sine wave in order to calculate the distribution of its ordinate. This latter distribution is of interest in the study of the scattered coherent reflected field from the sea surface [2].

Moreover we derive formulas for several integrals that are not readily available, and we exhibit some of the properties of the Kampé de Fériet functions associated with $K_{e_0}(a, z)$.

PRELIMINARY DEFINITIONS

The Pochhammer symbol $(a)_n$ is defined for nonnegative integers n as a ratio of gamma functions:

$$(a)_n \equiv \Gamma(a + n)/\Gamma(a) = a(a + 1) \ldots (a + n - 1)$$
$$(a)_0 \equiv 1 \tag{2}$$

Manuscript approved January 8, 1986.

Following Srivastava and Panda [3, p. 63] we define the Kampé de Fériet double hypergeometric functions:

$$F^{p:q;k}_{l:m;n}\left[\begin{matrix}(a_p):(b_q);(c_k);\\(\alpha_l):(\beta_m);(\gamma_n);\end{matrix}\; x, y\right] \equiv \sum_{r,s=0}^{\infty} \frac{\prod_{j=1}^{p}(a_j)_{r+s}\prod_{j=1}^{q}(b_j)_r\prod_{j=1}^{k}(c_j)_s}{\prod_{j=1}^{l}(\alpha_j)_{r+s}\prod_{j=1}^{m}(\beta_j)_r\prod_{j=1}^{n}(\gamma_j)_s}\frac{x^r}{r!}\frac{y^s}{s!}$$

where the Pochhamner symbols $(a)_n$ are defined by Eq. (2). For convergence

$$p+q<l+m+1,\ \ p+k<l+n+1,\ \ |x|<\infty,\ \ |y|<\infty, \text{ or}$$

$$p+q=l+m+1,\ \ p+k=l+n+1, \text{ and}$$

$$\begin{cases} |x|^{1/(p-l)}+|y|^{1/(p-l)}<1 & p>l \\ \max\{|x|,|y|\}<1 & p\leqslant l \end{cases}$$

As special cases we define

$$L[\alpha,\beta;\gamma,\delta;x,y] \equiv \sum_{m,n=0}^{\infty}\frac{(\alpha)_m(\beta)_n}{(\gamma)_{m+n}(\delta)_{m+n}}\frac{x^m}{m!}\frac{y^n}{n!} \qquad |x|<\infty,\ |y|<\infty \tag{3}$$

$$M[\alpha,\beta;\gamma,\delta;x,y] \equiv \sum_{m,n=0}^{\infty}\frac{(\alpha)_{m+n}(\beta)_n}{(\gamma)_{m+n}(\delta)_m}\frac{x^m}{m!}\frac{y^n}{n!} \qquad |x|<\infty,\ |y|<1 \tag{4}$$

We may then write

$$L[\alpha,\beta;\gamma,\delta;x,y] = F^{0:1;1}_{2:0;0}\left[\begin{matrix}-\ :\alpha;\beta;\\ \gamma,\delta:-;-;\end{matrix}\; x,y\right]$$

$$M[\alpha,\beta;\gamma,\delta;x,y] = F^{1:0;1}_{1:1;0}\left[\begin{matrix}\alpha:-;\beta;\\ \gamma:\delta;-;\end{matrix}\; x,y\right]$$

SOME ELEMENTARY PROPERTIES OF $M[\alpha,\beta;\gamma,\delta;x,y]$

Substituting [4, p. 266]

$$\frac{(\alpha)_p}{(\gamma)_p} = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\,\Gamma(\gamma-\alpha)}\int_0^1 t^{p+\alpha-1}(1-t)^{\gamma-\alpha-1}dt$$

UNCLASSIFIED

where Re $\gamma >$ Re $\alpha > 0$, and $p = m + n$ into Eq. (4), we deduce an integral representation for M:

$$M[\alpha, \beta; \gamma, \delta; x, y] = \frac{\Gamma(\gamma)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} \int_0^1 {}_0F_1[-; \delta; xt] t^{\alpha-1}(1-t)^{\gamma-\alpha-1}(1-yt)^{-\beta} dt$$

$$= \frac{\Gamma(\gamma)\Gamma(\delta)}{\Gamma(\alpha)\Gamma(\gamma-\alpha)} x^{-\frac{\delta-1}{2}} \int_0^1 I_{\delta-1}(2\sqrt{xt}) t^{\alpha-\frac{\delta+1}{2}} (1-t)^{\gamma-\alpha-1} (1-yt)^{-\beta} dt$$

Here we have used the equation

$$I_\nu(z) = \frac{(z/2)^\nu}{\Gamma(\nu+1)} {}_0F_1[-; \nu+1; z^2/4] \tag{5}$$

We obtain directly from Eq. (4) the generating relation

$$M[\alpha, \beta; \gamma, \delta; x, y] = \sum_{n=0}^{\infty} \frac{(\alpha)_n}{(\gamma)_n(\delta)_n} \frac{x^n}{n!} {}_2F_1[n+\alpha, \beta; n+\gamma; y] \tag{6}$$

We now prove the following

THEOREM: Suppose $-1 < \mathrm{Re}(\gamma - \alpha - \beta) < 0$, $|\arg y| < \pi$, $|\arg(1-y)| < \pi$. Then for $y \to 1$,

$$M[\alpha, \beta; \gamma, \delta; x, y] = \frac{\Gamma(\gamma)\,\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} {}_1F_2[\alpha; \gamma-\beta, \delta; x]$$

$$+ \frac{\Gamma(\gamma)\,\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\,\Gamma(\beta)} (1-y)^{\gamma-\alpha-\beta} {}_0F_1[-; \delta; x] + O(1-y) \tag{7}$$

or

$$M[\alpha, \beta; \gamma, \delta; x, y] = \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} {}_1F_2[\alpha; \gamma-\beta, \delta; x] + O(1-y)$$

$$+ \Gamma(\alpha+\beta-\gamma) \frac{\Gamma(\gamma)\Gamma(\delta)}{\Gamma(\alpha)\Gamma(\beta)} x^{-\frac{\delta-1}{2}} (1-y)^{\gamma-\alpha-\beta} I_{\delta-1}(2\sqrt{x}) \tag{8}$$

Proof: The following result is found in [4, Eq. (9.5.7), p. 249]: for $\alpha + \beta - \gamma \neq 0, \pm 1, \pm 2, \ldots$, $|\arg z| < \pi$, $|\arg(1-z)| < \pi$

$${}_2F_1[\alpha, \beta; \gamma; z] = \frac{\Gamma(\gamma)\,\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\,\Gamma(\gamma-\beta)} {}_2F_1[\alpha, \beta; 1+\alpha+\beta-\gamma; 1-z]$$

$$+ (1-z)^{\gamma-\alpha-\beta} \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)} {}_2F_1[\gamma-\alpha, \gamma-\beta; 1-\alpha-\beta+\gamma; 1-z]$$

Hence

$$
{}_2F_1[n+\alpha,\beta;n+\gamma;y]=\frac{\Gamma(n+\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(n+\gamma-\beta)}\,{}_2F_1[n+\alpha,\beta;1+\alpha+\beta-\gamma;1-y]
$$

$$
+(1-y)^{\gamma-\alpha-\beta}\frac{\Gamma(n+\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(n+\alpha)\Gamma(\beta)}\,{}_2F_1[\gamma-\alpha,n+\gamma-\beta;1-\alpha-\beta+\gamma;1-y]
$$

Now suppose that $-1 < \mathrm{Re}(\gamma-\alpha-\beta) < 0$. Then for $y \to 1$ we have

$$
{}_2F_1[n+\alpha,\beta;n+\gamma;y]=\frac{\Gamma(n+\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(n+\gamma-\beta)}+\frac{\Gamma(n+\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(n+\alpha)\Gamma(\beta)}(1-y)^{\gamma-\alpha-\beta}+O(1-y)
$$

$$
=\frac{(\gamma)_n}{(\gamma-\beta)_n}\,\frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\beta)\Gamma(\gamma-\alpha)}+\frac{(\gamma)_n}{(\alpha)_n}\,\frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)}\,(1-y)^{\gamma-\alpha-\beta}+O(1-y)
$$

Substituting this result into Eq. (6) gives

$$
M[\alpha,\beta;\gamma,\delta;x,y]=\frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)}\sum_{n=0}^{\infty}\frac{(\alpha)_n}{(\gamma-\beta)_n(\delta)_n}\frac{x^n}{n!}
$$

$$
+\frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)}(1-y)^{\gamma-\alpha-\beta}\sum_{n=0}^{\infty}\frac{x^n}{(\delta)_n n!}+O(1-y)
$$

from which we obtain Eq. (7). Then using Eq. (5) we obtain Eq. (8).

Employing series rearrangement we deduce

$$
M[\alpha,\beta;\gamma,\delta;x,tx]=\sum_{p=0}^{\infty}\frac{(\alpha)_p(\beta)_p}{(\gamma)_p}\frac{x^p}{p!}\,t^p\,{}_1F_2[-p;\delta,1-\beta-p;1/t] \tag{9}
$$

Using a general result of Srivastava [3, Eq. (30), p. 145] we find Eq. (9) in a different form, *viz*,

$$
M[\alpha,\beta;\gamma,\delta;tx,t]=\sum_{p=0}^{\infty}\frac{(\alpha)_p(\beta)_p}{(\gamma)_p}\frac{t^p}{p!}\,{}_1F_2[-p;\delta,1-\beta-p;x]
$$

From Eq. (9) it follows that

$$
M[\alpha,\beta;\gamma,\delta;x,tx]=\sum_{p=0}^{\infty}\frac{(\alpha)_p}{(\gamma)_p(\delta)_p}\frac{x^p}{p!}\,{}_3F_0[\beta,-p,1-\delta-p;-;t] \tag{10}
$$

Equation (10) may be obtained directly from [3, Eq. (60.ii), p. 194].

UNCLASSIFIED

We remark that it may be shown that $M[\alpha, 1; \gamma, \delta; x, y]$ converges on the unit circle $|y| = 1$ if and only if ${}_2F_1[\alpha, 1; \gamma; y]$ converges on $|y| = 1$.

SOME ELEMENTARY PROPERTIES OF $L[\alpha, \beta; \gamma, \delta; x, y]$

Using series rearrangement we find

$$L[\alpha, \beta; \gamma, \delta; x, tx] = \sum_{p=0}^{\infty} \frac{(\alpha)_p}{(\gamma)_p(\delta)_p} \frac{x^p}{p!} {}_2F_1[\beta, -p; 1 - \alpha - p; t]$$

This can also be obtained from [3, Eq. (30), p. 145] in a different form. Using Vandermonde's theorem [5, Eq. (1.7.7), p. 28]

$${}_2F_1[a, -p; c; 1] = (c - a)_p/(c)_p$$

we find

$${}_2F_1[\beta, -p; 1 - \alpha - p; 1] = \frac{(1 - \alpha - \beta - p)_p}{(1 - \alpha - p)_p} = \frac{(\alpha + \beta)_p}{(\alpha)_p}$$

so that we have a reduction formula for L, *viz*,

$$L[\alpha, \beta; \gamma, \delta; x, x] = \sum_{p=0}^{\infty} \frac{(\alpha + \beta)_p}{(\gamma)_p(\delta)_p} \frac{x^p}{p!} = {}_1F_2[\alpha + \beta; \gamma, \delta; x] \tag{11}$$

This result can be obtained also by using the following general result of Srivastava [3, Eq. (20), p. 55] applied to Eq. (3):

$$\sum_{m,n=0}^{\infty} c_{m+n}(\rho)_m(\sigma)_n \frac{x^{m+n}}{m!n!} = \sum_{n=0}^{\infty} c_n(\rho + \sigma)_n \frac{x^n}{n!}$$

provided each series is absolutely convergent.

We obtain directly from Eq. (3) the generating relation

$$L[\alpha, \beta; \gamma, \delta; x, y] = \sum_{m=0}^{\infty} \frac{(\alpha)_m}{(\gamma)_m(\delta)_m} \frac{x^m}{m!} {}_1F_2[\beta; m + \gamma, m + \delta; y] \tag{12}$$

Finally, using [3, Eq. (43), p. 150] we obtain

$$L[\alpha, \beta; \gamma, \delta; -x, x \tan^2\theta] = (\cos^2\theta)^\beta \sum_{n=0}^{\infty} \frac{(\beta)_n(\sin^2\theta)^n}{n!} {}_1F_2[\alpha - n; \gamma, \delta; -x]$$

A CLOSED FORM FOR $K_{e_0}(a, z)$

From Eq. (1) we write

$$K_{e_0}(\alpha/\beta, \beta) = \beta \int_0^1 e^{\alpha t} K_0(\beta t)\, dt \tag{13}$$

Using [6, p. 89] we find the following formulas:

$$\int_0^1 s^m K_0(zs)\, ds = \frac{K_0(z)}{m+1}\, {}_1F_2\left[1;\ \frac{m+1}{2},\ \frac{m+3}{2};\ \frac{z^2}{4}\right]$$

$$+ \frac{z K_1(z)}{(m+1)^2}\, {}_1F_2\left[1;\ \frac{m+3}{2},\ \frac{m+3}{2};\ \frac{z^2}{4}\right] \qquad m = 0,2,4,\ldots \tag{14}$$

$$\int_0^1 s^m K_0(zs)\, ds = \frac{2^{m-1}\,\Gamma\left(\frac{m+1}{2}\right)\Gamma\left(\frac{m+1}{2}\right)}{z^{m+1}}$$

$$- \frac{(m-1) K_0(z)}{z^2}\, {}_3F_0\left[1,\ \frac{1-m}{2},\ \frac{3-m}{2};\ -;\ \frac{4}{z^2}\right]$$

$$- \frac{K_1(z)}{z}\, {}_3F_0\left[1,\ \frac{1-m}{2},\ \frac{1-m}{2};\ -;\ \frac{4}{z^2}\right] \qquad m = 1,3,5,\ldots \tag{15}$$

Integrating term by term we find

$$\int_0^1 \exp(\alpha t) K_0(\beta t)\, dt = \int_0^1 \sum_{n=0}^{\infty} \frac{\alpha^n t^n}{n!} K_0(\beta t)\, dt = \sum_{n=0}^{\infty} \frac{\alpha^n}{n!} \int_0^1 t^n K_0(\beta t)\, dt$$

$$= \sum_{n=0,2,4,\ldots} \frac{\alpha^n}{n!} \int_0^1 t^n K_0(\beta t)\, dt + \sum_{n=1,3,5,\ldots} \frac{\alpha^n}{n!} \int_0^1 t^n K_0(\beta t)\, dt$$

$$= \sum_{n=0}^{\infty} \frac{\alpha^{2n}}{(2n)!} \int_0^1 t^{2n} K_0(\beta t)\, dt + \sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!} \int_0^1 t^{2n+1} K_0(\beta t)\, dt$$

UNCLASSIFIED

so that using Eqs. (14) and (15)

$$\int_0^1 \exp(\alpha t)\, K_0(\beta t)\, dt = \sum_{n=0}^{\infty} \frac{\alpha^{2n}}{(2n)!} \frac{K_0(\beta)}{2n+1}\, {}_1F_2\left[1;\ \frac{2n+1}{2},\ \frac{2n+3}{2};\ \frac{\beta^2}{4}\right]$$

$$+ \sum_{n=0}^{\infty} \frac{\alpha^{2n}}{(2n)!} \frac{\beta K_1(\beta)}{(2n+1)^2}\, {}_1F_2\left[1;\ \frac{2n+3}{2},\ \frac{2n+3}{2};\ \frac{\beta^2}{4}\right]$$

$$+ \sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!}\, 2^{2n}\, \frac{\Gamma(n+1)\Gamma(n+1)}{\beta^{2n+2}}$$

$$- \sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!}\, (2n)\, \frac{K_0(\beta)}{\beta^2}\, {}_3F_0[1, -n, 1-n; -; 4/\beta^2]$$

$$- \sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!} \frac{K_1(\beta)}{\beta}\, {}_3F_0[1, -n, -n; -; 4/\beta^2] \tag{16}$$

We shall consider each of the above five sums in the order in which they appear. We find

$$\sum_{n=0}^{\infty} \frac{\alpha^{2n}}{(2n+1)!}\, {}_1F_2[1;\ n+1/2,\ n+3/2;\ \beta^2/4]$$

$$= \sum_{n=0}^{\infty} \frac{1}{(3/2)_n} \frac{(\alpha^2/4)^n}{n!}\, {}_1F_2[1;\ n+1/2,\ n+3/2;\ \beta^2/4] = L\ [1/2,\ 1;\ 1/2,\ 3/2;\ \alpha^2/4,\ \beta^2/4];$$

$$\sum_{n=0}^{\infty} \frac{\alpha^{2n}}{(2n+1)(2n+1)!}\, {}_1F_2[1;\ n+3/2,\ n+3/2;\ \beta^2/4]$$

$$= \sum_{n=0}^{\infty} \frac{(1/2)_n}{(3/2)_n (3/2)_n} \frac{(\alpha^2/4)^n}{n!}\, {}_1F_2[1;\ n+3/2,\ n+3/2;\ \beta^2/4] = L\ [1/2,\ 1;\ 3/2,\ 3/2;\ \alpha^2/4,\ \beta^2/4]$$

where in the latter two cases we have used Eq. (12);

$$\sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!}\, 2^{2n}\, \frac{\Gamma(n+1)\Gamma(n+1)}{\beta^{2n+2}} = \frac{\sin^{-1}(\alpha/\beta)}{\sqrt{\beta^2 - \alpha^2}} \qquad |\alpha/\beta| \leqslant 1, \quad \alpha \neq \pm\beta$$

where we have used [9, Eq. (9.121-14), p. 1041] the result ${}_2F_1[1, 1; 3/2; \sin^2 z] = z/\sin z \cos z$;

$$\sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!} (2n)\, {}_3F_0[1, -n, 1-n; -; 4/\beta^2] = \sum_{n=0}^{\infty} \frac{\alpha^{2n+3}}{(2n+3)!} (2n+2)\, {}_3F_0[1, -1-n, -n; -; 4/\beta^2]$$

$$= \frac{\alpha^3}{3} \sum_{n=0}^{\infty} \frac{1}{(5/2)_n} \frac{(\alpha^2/4)^n}{n!}\, {}_3F_0[1, -1-n, -n; -; 4/\beta^2] = \frac{\alpha^3}{3} M\left[2, 1; \frac{5}{2}, 2; \frac{\alpha^2}{4}, \frac{\alpha^2}{\beta^2}\right];$$

and finally

$$\sum_{n=0}^{\infty} \frac{\alpha^{2n+1}}{(2n+1)!}\, {}_3F_0\, [1, -n, -n; -; 4/\beta^2]$$

$$= \alpha \sum_{n=0}^{\infty} \frac{1}{(3/2)_n} \frac{(\alpha^2/4)^n}{n!}\, {}_3F_0[1, -n, -n; -; 4/\beta^2] = \alpha\, M\left[1, 1; \frac{3}{2}, 1; \frac{\alpha^2}{4}, \frac{\alpha^2}{\beta^2}\right]$$

where in the latter two cases we have used Eq. (10).

Defining

$$L_0(x, y) \equiv \sum_{m,n=0}^{\infty} \frac{(1/2)_m (1)_n}{(3/2)_{m+n} (3/2)_{m+n}} \frac{x^m}{m!} \frac{y^n}{n!} = L[1/2, 1; 3/2, 3/2; x, y]$$

$$L_1(x, y) \equiv \sum_{m,n=0}^{\infty} \frac{(1/2)_m (1)_n}{(1/2)_{m+n} (3/2)_{m+n}} \frac{x^m}{m!} \frac{y^n}{n!} = L\, [1/2, 1; 1/2, 3/2; x, y]$$

$$M_0(x, y) \equiv \sum_{m,n=0}^{\infty} \frac{(1)_{m+n}}{(3/2)_{m+n}} \frac{(1)_n}{(1)_m} \frac{x^m}{m!} \frac{y^n}{n!} = M\, [1, 1; 3/2, 1; x, y]$$

$$M_1(x, y) \equiv \sum_{m,n=0}^{\infty} \frac{(2)_{m+n}}{(5/2)_{m+n}} \frac{(1)_n}{(2)_m} \frac{x^m}{m!} \frac{y^n}{n!} = M[2, 1; 5/2, 2; x, y]$$

we have from Eq. (16) and the above results

$$\int_0^1 \exp\,(\alpha t) K_0(\beta t)\, dt = K_1(\beta)\left[\beta L_0(\alpha^2/4, \beta^2/4) - \frac{\alpha}{\beta} M_0(\alpha^2/4, \alpha^2/\beta^2)\right]$$

$$+ K_0(\beta)\left[L_1(\alpha^2/4, \beta^2/4) - \frac{\alpha^3}{3\beta^2} M_1(\alpha^2/4, \alpha^2/\beta^2)\right] + \frac{\sin^{-1}(\alpha/\beta)}{\sqrt{\beta^2 - \alpha^2}} \qquad (17)$$

UNCLASSIFIED

which we may write using Eq. (13)

$$K_{e_0}(a, z) = z\, K_1(z) \left[z\, L_0 \left(\frac{a^2 z^2}{4}, \frac{z^2}{4} \right) - a\, M_0 \left(\frac{a^2 z^2}{4}, a^2 \right) \right]$$

$$+ z\, K_0(z) \left[L_1 \left(\frac{a^2 z^2}{4}, \frac{z^2}{4} \right) - \frac{a^3 z}{3} M_1 \left(\frac{a^2 z^2}{4}, a^2 \right) \right] + \frac{\sin^{-1} a}{\sqrt{1 - a^2}} \tag{18}$$

We have then given $K_{e_0}(a, z)$ in terms of elementary, MacDonald, and Kampé de Fériet functions.

We remark that in view of Eq. (6) and the definitions of M_0 and M_1

$$M_0(x, y) = \sum_{n=0}^{\infty} \frac{1}{(3/2)_n} \frac{x^n}{n!}\, {}_2F_1[1, n + 1; n + 3/2; y]$$

$$M_1(x, y) = \sum_{n=0}^{\infty} \frac{1}{(5/2)_n} \frac{x^n}{n!}\, {}_2F_1[1, n + 2; n + 5/2; y]$$

Since each of the Gauss hypergeometric functions above is conditionally convergent on the unit circle $|y| = 1$ except at $y = 1$ we see that $M_0(x, y)$ and $M_1(x, y)$ are conditionally convergent on $|y| = 1$ except at $y = 1$.* Hence Eq. (17) is valid for $|\alpha/\beta| \leqslant 1$, $\alpha \neq \pm\beta$ and Eq. (18) is valid for $|a| \leqslant 1$, $a \neq \pm 1$. We shall show shortly that Eq. (17) is valid in the limit even when $\alpha = \pm\, \beta$. See Ref. 1 for other representations of $K_{e_0}(a, z)$.

In a future report (Part II) it shall be shown that Eq. (18) is easily extended to the entire complex a-plane in terms of elementary, MacDonald and Kampé de Fériet functions.

KING'S INTEGRAL

Using properties of L and M we have derived earlier we shall derive (a formula for) King's integral [6, Eq. (12), p. 123]:

$$\int_0^{\alpha} \exp t\, K_0(t)\, dt = \alpha \exp \alpha [K_0(\alpha) + K_1(\alpha)] - 1 \tag{19}$$

that is we shall show Eq. (17) is valid in the limit for $\alpha = \beta$. Using Eq. (8) we find for $\alpha \to \beta$

$$M_0(\alpha^2/4, \alpha^2/\beta^2) = \frac{\pi}{2} \frac{I_0(\alpha)}{\sqrt{1 - \alpha^2/\beta^2}} - \cosh \alpha + O(1 - \alpha^2/\beta^2)$$

$$M_1(\alpha^2/4, \alpha^2/\beta^2) = \frac{3}{\alpha} \left[\frac{\pi}{2} \frac{I_1(\alpha)}{\sqrt{1 - \alpha^2/\beta^2}} - \sinh \alpha + O(1 - \alpha^2/\beta^2) \right]$$

*Also see remark on top of p. 5.

Substituting these equations into Eq. (17) gives

$$\int_0^1 \exp(\alpha t) K_0(\beta t)\, dt = \frac{\sin^{-1}(\alpha/\beta) - \frac{\pi}{2}[\alpha K_1(\beta)\, I_0(\alpha) + (\alpha^2/\beta)\, K_0(\beta)\, I_1(\alpha)]}{\sqrt{\beta^2 - \alpha^2}} + \frac{\alpha}{\beta} K_1(\beta) \cosh \alpha$$

$$+ \frac{\alpha^2}{\beta^2} K_0(\beta) \sinh \alpha + K_0(\beta) L_1(\alpha^2/4, \beta^2/4) + \beta K_1(\beta)\, L_0(\alpha^2/4, \beta^2/4) + O(1 - \alpha^2/\beta^2) \tag{20}$$

Using the reduction formula Eq. (11) for L we deduce

$$L_0(x^2/4,\ x^2/4) = \frac{\sinh x}{x}$$

$$L_1(x^2/4,\ x^2/4) = \cosh x$$

Now holding β fixed and letting $\alpha \to \beta$ we obtain after simplification

$$\int_0^1 \exp(\beta t) K_0(\beta t)\, dt = [K_0(\beta) + K_1(\beta)] \exp \beta + \lim_{\alpha \to \beta} J(\alpha, \beta)$$

where $J(\alpha, \beta)$ is the first term on the right-hand side of Eq. (20). We find however that

$$\lim_{\alpha \to \beta} [\text{numerator } J(\alpha, \beta)] = \frac{\pi}{2}[1 - \beta K_1(\beta)\, I_0(\beta) - \beta K_0(\beta)\, I_1(\beta)] = 0$$

$$\lim_{\alpha \to \beta} [\text{denominator } J(\alpha, \beta)] = 0$$

so that on applying L'hospital's rule we have

$$\lim_{\alpha \to \beta} J(\alpha, \beta) = -1/\beta$$

Hence

$$\int_0^1 \exp(\beta t) K_0(\beta t)\, dt = [K_0(\beta) + K_1(\beta)] \exp \beta - 1/\beta$$

and a simple transformation now gives Eq. (19). We may perform a similar analysis for $\alpha \to -\beta$ to obtain

$$\int_0^1 \exp(-\beta t) K_0(\beta t)\, dt = [K_0(\beta) - K_1(\beta)] \exp(-\beta) + 1/\beta$$

UNCLASSIFIED

A DISTRIBUTION FOR THE ELEVATION OF A SINE WAVE

Consider the random variable $y = H \sin \theta$, where H is a random variable with density $K(H, \epsilon)$, $|H| < \infty$, and θ is a random variable, independent of H, with density

$$U(\theta) = \pi^{-1} \qquad |\theta| \leqslant \pi/2$$
$$= 0 \qquad |\theta| > \pi/2$$

Let $D(y, \epsilon)$ be the density function for y. It is shown in Ref. 2 that

$$D(y, \epsilon) = \frac{1}{\pi} \int_{-\infty}^{-|y|} \frac{K(H, \epsilon)\, dH}{\sqrt{H^2 - y^2}} + \frac{1}{\pi} \int_{|y|}^{\infty} \frac{K(H, \epsilon)\, dH}{\sqrt{H^2 - y^2}} \tag{21}$$

Rice [7] and Cartwright and Longuet-Higgins [8] have derived an expression for the statistical distribution of the maxima of a random function that may be expressed in the form

$$K(H, \epsilon) = \frac{\epsilon}{\sigma_H \sqrt{2\pi}} \exp\left(\frac{-H^2}{2\epsilon^2 \sigma_H^2}\right) + \frac{\sqrt{1-\epsilon^2}}{2\sigma_H^2} H \exp\left(\frac{-H^2}{2\sigma_H^2}\right)\left[1 + \operatorname{erf}\left(\frac{\sqrt{2}}{2} \frac{H}{\sigma_H} \frac{\sqrt{1-\epsilon^2}}{\epsilon}\right)\right] \tag{22}$$

Here σ_H is the standard deviation of H, and $0 < \epsilon < 1$ is known as the spectral width parameter. It is shown in Ref. 2 that the standard deviation σ of y is given by

$$\sigma = \sigma_H / (\sqrt{2}\eta)$$

where η is defined by

$$\eta \equiv \left[1 + \frac{\pi}{2}(1 - \epsilon^2)\right]^{-1/2}$$

Substituting Eq. (22) into Eq. (21) and using the latter result gives

$$D(y, \epsilon) = \frac{\epsilon}{2\pi^{3/2}\eta\sigma} \exp\left(\frac{-y^2}{8\epsilon^2\eta^2\sigma^2}\right) K_0\left(\frac{y^2}{8\epsilon^2\eta^2\sigma^2}\right) + \frac{\sqrt{1-\epsilon^2}}{\pi\eta\sigma} \exp\left(\frac{-y^2}{4\eta^2\sigma^2}\right) \Psi\left(\frac{\sqrt{1-\epsilon^2}}{\epsilon}, \frac{y}{2\eta\sigma}\right) \tag{23}$$

where the function $\Psi(k, u)$ is defined by

$$\Psi(k, u) \equiv \int_0^{\infty} \exp(-s^2) \operatorname{erf}(k\sqrt{u^2 + s^2})\, ds \tag{24}$$

For real u and k it is shown in Ref. 2 that

$$\pi^{1/2} \int_0^{\infty} \exp(-s^2) \operatorname{erf}(k\sqrt{u^2 + s^2})\, ds = \tan^{-1} k + \frac{k}{1 + k^2} \int_0^{\frac{1}{2}u^2(1+k^2)} \exp\left(\frac{1 - k^2}{1 + k^2} s\right) K_0(s)\, ds$$

Using Eqs. (1) and (24) this may be written

$$\Psi(k, u) = \frac{\tan^{-1}(k)}{\pi^{1/2}} + \frac{1}{\pi^{1/2}} \frac{k}{1+k^2} K_{e_0}\left[\frac{1-k^2}{1+k^2}, \frac{1}{2}u^2(1+k^2)\right]$$

We may then write Eq. (23)

$$D(y, \epsilon) = \frac{\epsilon}{2\pi^{3/2}\eta\sigma} \exp\left[\frac{-y^2}{8\epsilon^2\eta^2\sigma^2}\right] K_0\left[\frac{y^2}{8\epsilon^2\eta^2\sigma^2}\right]$$

$$+ \frac{\sqrt{1-\epsilon^2}}{\pi^{3/2}\eta\sigma} \exp\left[\frac{-y^2}{4\eta^2\sigma^2}\right] [\cos^{-1}\epsilon + \epsilon\sqrt{1-\epsilon^2}\, K_{e_0}(2\epsilon^2 - 1, y^2/8\epsilon^2\eta^2\sigma^2)]$$

where $K_{e_0}(a, z)$ is given by Eq. (18).

SOME INTEGRALS RELATED TO $K_{e_0}(a, z)$

The following integrals can easily be obtained from Eq. (17):

$$\int_0^1 \sin(\alpha t) K_0(\beta t)\, dt = \frac{\sinh^{-1}(\alpha/\beta)}{\sqrt{\alpha^2+\beta^2}} - \frac{\alpha}{\beta} K_1(\beta)\, M_0(-\alpha^2/4, -\alpha^2/\beta^2)$$

$$+ \frac{\alpha^3}{3\beta^2} K_0(\beta)\, M_1(-\alpha^2/4, -\alpha^2/\beta^2) \qquad |\alpha/\beta| \leqslant 1, \quad \alpha \neq \pm i\beta$$

$$\int_0^1 \cos(\alpha t)\, K_0(\beta t)\, dt = \beta K_1(\beta)\, L_0(-\alpha^2/4, \beta^2/4) + K_0(\beta) L_1(-\alpha^2/4, \beta^2/4)$$

Further, using the result [2] for $0 < |\alpha| < 1$, $0 < x$

$$K_{e_0}(\alpha, x) = \operatorname{sgn}\alpha \left\{\exp(\alpha x)\left[\int_0^\infty \frac{\cos(\alpha x t)\, dt}{(1+t^2)\sqrt{1+\alpha^2 t^2}} + \int_0^\infty \frac{t \sin(\alpha x t)\, dt}{(1+t^2)\sqrt{1+\alpha^2 t^2}}\right] - \frac{\cos^{-1}(|\alpha|)}{\sqrt{1-\alpha^2}}\right\}$$

we find for $0 < \alpha < \beta$

$$\int_0^\infty \frac{\cos(\alpha x)\, dx}{(1+x^2)\sqrt{\beta^2+\alpha^2 x^2}} = \cosh\alpha \left\{\frac{\pi/2}{\sqrt{\beta^2-\alpha^2}} - \frac{\alpha}{\beta} K_1(\beta)\, M_0(\alpha^2/4, \alpha^2/\beta^2) - \frac{\alpha^3}{3\beta^2} K_0(\beta)\, M_1(\alpha^2/4, \alpha^2/\beta^2)\right\}$$

$$- \sinh\alpha \left\{\beta K_1(\beta) L_0(\alpha^2/4, \beta^2/4) + K_0(\beta) L_1(\alpha^2/4, \beta^2/4)\right\}$$

UNCLASSIFIED

$$\int_0^\infty \frac{x \sin(\alpha x)\, dx}{(1+x^2)\sqrt{\beta^2+\alpha^2 x^2}} = \cosh\alpha \left\{\beta K_1(\beta) L_0(\alpha^2/4, \beta^2/4) + K_0(\beta)\, L_1(\alpha^2/4, \beta^2/4)\right\}$$

$$- \sinh\alpha \left\{\frac{\pi/2}{\sqrt{\beta^2-\alpha^2}} - \frac{\alpha}{\beta} K_1(\beta) M_0(\alpha^2/4, \alpha^2/\beta^2) - \frac{\alpha^3}{3\beta^2} K_0(\beta)\, M_1(\alpha^2/4, \alpha^2/\beta^2)\right\}$$

In addition [9, Eq. (3.367), p. 316] we have

$$\int_0^\infty \frac{e^{-pt} \sin\theta\, dt}{(1+t+\cos\theta)\sqrt{t^2+2t}} = \exp\left[2p\cos^2\frac{\theta}{2}\right] [\theta - \sin\theta\, K_{e_0}(-\cos\theta, p)] \qquad \mathrm{Re}\, p > 0$$

CONCLUSIONS

The Kampé de Fériet functions have been used to put in closed form the incomplete Lipschitz-Hankel integral $K_{e_0}(a, z)$ and several related integrals that are not readily available and are of interest in mathematical physics and applications. Some of the properties of the Kampé de Fériet functions associated with $K_{e_0}(a, z)$ are derived. These properties are useful in deriving additional results quickly. As an example we have given an elementary derivation of a closed form for King's integral based on generating function techniques.

In addition, the utility of a closed form for $K_{e_0}(a, z)$ is indicated by deriving a certain density function that is associated with the scattered coherent return from the sea surface.

REFERENCES

1. M.M. Agrest and M.S. Maksimov, *Theory of Incomplete Cylindrical Functions and Their Applications*, Springer-Verlag, 1971.

2. A.R. Miller and E. Vegh, "A Family of Curves for the Rough Surface Reflection Coefficient," NRL Report 8898, July 1985.

3. H.M. Srivastava and H.L. Manocha, *A Treatise on Generating Functions*, Ellis Horwood Limited, 1984.

4. N.N. Lebedev, *Special Functions and Their Applications*, Dover, 1972.

5. L.J. Slater, *Generalized Hypergeometric Functions*, Cambridge Univ. Press, 1966.

6. Y.L. Luke, *Integrals of Bessel Functions*, McGraw-Hill, 1962.

7. S.O. Rice, "Mathematical Analysis of Random Noise," *Bell System Tech. J.* **24**, 46, 1945.

8. D.E. Cartwright and M.S. Longuet-Higgins, "The Statistical Distribution of the Maxima of a Random Function," *Proc. Royal Soc. London*, Series A, **237**, 212-232, 1956.

9. I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, 1980.