

An Analytical Necessary and Sufficient Condition for a Riemannian Manifold to be Complete

WILLIAM B. GORDON

Mathematics Research Center
Report 72-3

Mathematics and Information Sciences Division

July 6, 1972



NAVAL RESEARCH LABORATORY
Washington, D.C.

An Analytical Necessary and Sufficient Condition for a Riemannian Manifold to be Complete

WILLIAM B. GORDON

*Mathematics Research Center
Mathematics and Information Sciences Division*

Abstract: If M is a (not necessarily complete) riemannian manifold with metric tensor g_{ij} and f is any proper real-valued function on M , then M is necessarily complete with respect to the metric $\tilde{g}_{ij} = g_{ij} + (\partial f / \partial x^i)(\partial f / \partial x^j)$. Using this construction one can easily prove that a riemannian manifold is complete if and only if it supports a proper function whose gradient is bounded in modulus.

1. It is known that every differentiable manifold supports a complete riemannian structure. This is a consequence of Whitney's Embedding Theorem. (See the concluding remark at the end of Section 4.) Moreover, Nomizu and Ozeki [1] have shown that every riemannian manifold is conformally equivalent to a complete riemannian manifold. In this report we shall describe a method for constructing complete riemannian metrics which is exceedingly simple and provides a necessary and sufficient condition for the completeness of a riemannian structure. Namely, we shall show that a riemannian manifold is complete if and only if it supports a proper function whose gradient is bounded in modulus.

We begin by recalling some basic definitions. All manifolds under discussion will be assumed to be connected paracompact Hausdorff spaces which are locally homeomorphic to open euclidean m -discs. A riemannian manifold M of class C^3 with metric tensor $g = (g_{ij})$ is said to be *complete* if either of the following two equivalent conditions hold:

- (i) Every geodesic arc $\gamma = \gamma(t)$ can be extended for all values of t .
- (ii) M is complete with respect to the distance function $d(p, q) =$ (infimum of the set of arc lengths of paths joining p to q).

Recall that in order for M to be complete it suffices that there exist a single point p such that every geodesic $\gamma = \gamma(t)$ with $\gamma(0) = p$ can be extended for all values of t ; also, every pair of points of a complete riemannian manifold can be joined by a geodesic of minimum length. (For details and references see e.g. [2] or [3].) If $b = (b_{ij})$ is a symmetric tensor field; we write $(b_{ij}) > 0$ to indicate that the form $(b_{ij}(p))$ is positive definite at every point p of M , i.e., that $\sum b_{ij}(p)\xi^i\xi^j > 0$ for every p in M and every nonzero vector ξ in $T_p(M)$. By definition, a riemannian metric is a symmetric tensor field $\gamma = (\gamma_{ij})$ of class C^2 which is positive definite at every point. Finally, we recall that a continuous map f is said to be *proper* if $f^{-1}(K)$ is compact whenever K is compact. Our results are contained in the following two theorems.

NRL Problem B01-11; Project RR 003-02-41-6153. This is a final report on one phase of the problem; work is continuing on other phases. Manuscript submitted May 2, 1972.

THEOREM 1. *Let M be a (not necessarily complete) riemannian manifold of class C^3 with metric tensor $g = (g_{ij})$, and let f be any proper real-valued function on M of class C^3 . Then M is complete with respect to the metric $\tilde{g} = (g_{ij})$, where*

$$\tilde{g}_{ij} = g_{ij} + f_i f_j, \quad \left(f_i = \frac{\partial f}{\partial x^i} \right). \quad (1)$$

THEOREM 2. *Let M be as above. Then M is complete with respect to g if and only if there exists a proper C^3 function f on M such that*

$$(g_{ij} - f_i f_j) > 0. \quad (2)$$

Equivalently, a riemannian manifold is complete if and only if it supports a proper C^3 function f whose gradient is bounded in modulus, ($|\nabla f| \leq \text{constant}$).

Proof of Theorem 1.

2. In a few words, to prove the theorem we first observe that \tilde{g} is the metric that g induces on the graph of f , and then use the propriety of f to show that its graph is Cauchy complete.

Let the product manifold $M \times R$ be endowed with the usual product metric, so that the element of arc on $M \times R$ becomes $\overline{ds}^2 = \sum g_{ij} dx^i dx^j + \overline{dz}^2$, where $x = (x^1, \dots, x^m)$ represents a local coordinate chart on M and z denotes a general point of R . The *graph* of f will be defined to be the subspace of $M \times R$ which consists of all pairs $\{(p, f(p)) | p \in M\}$ and which is endowed with the obvious differentiable structure which makes G diffeomorphic to M . Hereafter we identify G with M . Now \tilde{g} is the metric which G inherits from its embedding in $M \times R$, and therefore the theorem will be proved once we show that every \tilde{g} -Cauchy sequence in G contains a \tilde{g} -convergent subsequence.

But if $\{(p_n, f(p_n))\}$ is a Cauchy sequence in G , then $\{f(p_n)\}$ is a Cauchy sequence in R because

$$d_G[(p, z), (p', z')] \geq d_{M \times R}[(p, z), (p', z')] \geq |z - z'|.$$

Therefore, by passing to a subsequence we can assume that $f(p_n) \rightarrow z$ for some z in R , so that $\{z, f(p_1), f(p_2), \dots\}$ is a compact subset of R . But $\{p_1, p_2, \dots\} \subset f^{-1}\{z, f(p_1), f(p_2), \dots\}$ and f is proper. Therefore the sequence $\{p_n\}$ contains a convergent subsequence since it is contained in a compact set.

Remark 1. The proof generalizes the following construction: Let A be a closed subset of R^2 and let f be a C^3 function defined on $R^2 - A$. Let G be the surface embedded in R^3 which lies over $R^2 - A$ and is given by the relation $z = f(x, y)$. Then G is complete (with respect to the induced metric) if f is a proper function (on $R^2 - A$).

Remark 2. Let g and \tilde{g} be two metric tensors on M . It is a trivial matter to verify that M is complete with respect to \tilde{g} if M is complete with respect to g and $(\tilde{g}_{ij} - g_{ij})$ is positive semi-definite at every point. Hence from (1) it follows that the graph of any function on a manifold is complete if the manifold is itself complete.

Proof of Theorem 2.

3. Let f be a proper C^3 function which satisfies (2). Then the assertion that M is complete is an immediate consequence of Theorem 1 and the relations $g_{ij} = (g_{ij} - f_i f_j) + f_i f_j$.

We shall raise and lower indices in the usual tensorial fashion and employ the tensor summation convention. If f satisfies (2), then $|\nabla f| < 1$; for multiplying (2) by $f^i f^j$ and summing on repeated indices, we get $|\nabla f|^2 - |\nabla f|^4 > 0$. On the other hand suppose $|\nabla f| \leq \text{constant}$. Multiplying f by a suitably small constant we can assume that $|\nabla f| < 1$. Then for any vector ξ we have

$$g_{ij} \xi^i \xi^j - (f_i \xi^i)(f_j \xi^j) = |\xi|^2 - (\nabla f, \xi)^2 \geq |\xi|^2(1 - |\nabla f|^2).$$

that is, $(g_{ij} - f_i f_j) > 0$, and we have shown that the existence of a proper function satisfying (2) is equivalent to the existence of a proper function whose gradient is bounded in modulus.

Finally, to complete the proof we have to show that every complete riemannian manifold M supports a proper C^3 function f whose gradient is bounded in modulus. By a well-known theorem of Nash we can assume the existence of an isometric embedding $j: M \rightarrow R^N$ for some N . Then M is complete if and only if M is a closed subspace of R^N if and only if j is proper. Let $w = (w^1, \dots, w^N)$ be standard euclidean coordinates on R^N , and let $F(w) = \log(1 + |w|^2)$. Then F is a proper function on R^N and $|\nabla F| \leq 1$. Let $f = F \circ j = F/M$. Then f is proper because the composition of proper maps is proper, and $|\nabla f| \leq 1$ because at each point p of M , $\nabla f(p)$ is the orthogonal projection of $\nabla F(p)$ onto $T_p(M)$, so that $|\nabla f(p)| \leq |\nabla F(p)|$.

4. We conclude by remarking that the theorem of Nomizu and Ozeki is an easy consequence of Theorem 2. In fact, we have the following corollary.

COROLLARY. *Let f be any proper C^3 function on M , and let $g = (g_{ij})$ be any metric tensor on M . Then M is complete with respect to the metric \tilde{g} , where*

$$\tilde{g}_{ij} = e^{|\nabla f|^2} g_{ij}.$$

Proof. Let $\nabla_g, \nabla_{\tilde{g}}, |\cdot|_g, |\cdot|_{\tilde{g}}$ denote the gradient operators and norms associated with g and \tilde{g} respectively. Then

$$\left| \nabla_{\tilde{g}} f \right|_{\tilde{g}}^2 = \left| \nabla_g f \right|_g^2 \exp\left(-\left| \nabla_g f \right|_g^2\right) \leq \frac{1}{e}.$$

Remark. The existence of a proper junction on M is guaranteed by Whitney's Embedding Theorem, which asserts that every manifold can be embedded as a *closed* submanifold of euclidean space. This being given, one then proceeds to construct a proper junction on M as was done at the end of the last section. Whitney's theorem also implies the existence of a complete metric on M . For if M is embedded as a closed submanifold of R^N one can define g to be the metric which is induced by the embedding.

REFERENCES

1. K. Nomizu and H. Ozeki, "The existence of complete Riemannian metrics," Proc. Am. Math. Soc. 12 (1961), 889-891.
2. R. Hermann, *Differential Geometry and the Calculus of Variations*, Academic Press, New York, 1968.
3. S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Interscience, New York, 1963.

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

| | | | |
|--|---|---|--|
| 1. ORIGINATING ACTIVITY <i>(Corporate author)</i> Naval Research Laboratory Washington, D.C. 20390 | | 2a. REPORT SECURITY CLASSIFICATION Unclassified | |
| 2b. GROUP | | | |
| 3. REPORT TITLE AN ANALYTICAL NECESSARY AND SUFFICIENT CONDITION FOR A RIEMANNIAN MANIFOLD TO BE COMPLETE | | | |
| 4. DESCRIPTIVE NOTES <i>(Type of report and inclusive dates)</i> This is a final report on one phase of the problem. Work is continuing on other phases. | | | |
| 5. AUTHOR(S) <i>(First name, middle initial, last name)</i> William B. Gordon | | | |
| 6. REPORT DATE July 6, 1972 | 7a. TOTAL NO. OF PAGES 6 | 7b. NO. OF REFS 3 | |
| 8a. CONTRACT OR GRANT NO. NRL Problem B01-11 | 9a. ORIGINATOR'S REPORT NUMBER(S) NRL Report 7441 | | |
| b. PROJECT NO. RR 003-02-41-6153 | 9b. OTHER REPORT NO(S) <i>(Any other numbers that may be assigned this report)</i> MRC Report 72-3 | | |
| c. | | | |
| d. | | | |
| 10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited. | | | |
| 11. SUPPLEMENTARY NOTES | | 12. SPONSORING MILITARY ACTIVITY Department of the Navy Office of Naval Research Arlington, Virginia 22217 | |
| 13. ABSTRACT If M is a (not necessarily complete) riemannian manifold with metric tensor g_{ij} and f is any proper real-valued function on M , then M is necessarily complete with respect to the metric $\tilde{g}_{ij} = g_{ij} + (\partial f / \partial x^i)(\partial f / \partial x^j)$. Using this construction one can easily prove that a riemannian manifold is complete if and only if it supports a proper function whose gradient is bounded in modulus. | | | |

| 14. KEY WORDS | LINK A | | LINK B | | LINK C | |
|------------------------------|--------|----|--------|----|--------|----|
| | ROLE | WT | ROLE | WT | ROLE | WT |
| Complete Riemannian Manifold | | | | | | |