

Naval Research Laboratory

Washington, DC 20375-5000



NRL 9146 COPY 1

NRL Report 9146

Reduction Formulae for Kampé de Fériet Functions $F_{q:1;0}^{p:2;1}$

ALLEN R. MILLER

*Industrial Engineering Services Branch
Engineering Services Division*

March 23, 1989

22-336



ENRICHMENT RESOURCE CENTER

Approved for public release; distribution unlimited.

SECURITY CLASSIFICATION OF THIS PAGE

| REPORT DOCUMENTATION PAGE | | | | Form Approved OMB No. 0704-0188 | | | | | | | | | | | | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|----------------------------------------|------------------------------------|--|--|--|--|--|--|--|--|--|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| 1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED | | 1b. RESTRICTIVE MARKINGS | | | | | | | | | | | | | | |
| 2a. SECURITY CLASSIFICATION AUTHORITY | | 3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited. | | | | | | | | | | | | | | |
| 2b. DECLASSIFICATION / DOWNGRADING SCHEDULE | | | | | | | | | | | | | | | | |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Report 9146 | | 5. MONITORING ORGANIZATION REPORT NUMBER(S) | | | | | | | | | | | | | | |
| 6a. NAME OF PERFORMING ORGANIZATION Naval Research Laboratory | 6b. OFFICE SYMBOL (<i>If applicable</i>) Code 2330 | 7a. NAME OF MONITORING ORGANIZATION | | | | | | | | | | | | | | |
| 6c. ADDRESS (<i>City, State, and ZIP Code</i>) Washington, DC 20375-5000 | | 7b. ADDRESS (<i>City, State, and ZIP Code</i>) | | | | | | | | | | | | | | |
| 8a. NAME OF FUNDING / SPONSORING ORGANIZATION | 8b. OFFICE SYMBOL (<i>If applicable</i>) | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER | | | | | | | | | | | | | | |
| 8c. ADDRESS (<i>City, State, and ZIP Code</i>) | | 10. SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. PROJECT NO. TASK NO. WORK UNIT ACCESSION NO. | | | | | | | | | | | | | | |
| 11. TITLE (<i>Include Security Classification</i>) Reduction Formulae for Kampé de Fériet Functions $F_{q:1;0}^{p:2;1}$ | | | | | | | | | | | | | | | | |
| 12. PERSONAL AUTHOR(S) Miller, A.R. | | | | | | | | | | | | | | | | |
| 13a. TYPE OF REPORT Final | 13b. TIME COVERED FROM _____ TO _____ | 14. DATE OF REPORT (<i>Year, Month, Day</i>) 1989 March 23 | 15. PAGE COUNT 15 | | | | | | | | | | | | | |
| 16. SUPPLEMENTARY NOTATION | | | | | | | | | | | | | | | | |
| 17. COSATI CODES <table border="1"> <tr><th>FIELD</th><th>GROUP</th><th>SUB-GROUP</th></tr> <tr><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td></tr> <tr><td></td><td></td><td></td></tr> </table> | | FIELD | GROUP | SUB-GROUP | | | | | | | | | | 18. SUBJECT TERMS (<i>Continue on reverse if necessary and identify by block number</i>) Kampé de Fériet functions Hypergeometric functions | | |
| FIELD | GROUP | SUB-GROUP | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | |
| 19. ABSTRACT (<i>Continue on reverse if necessary and identify by block number</i>) Several reduction formulae for Kampé de Fériet functions $F_{q:1;0}^{p:2;1}$ are given in terms of a finite number of generalized hypergeometric functions. | | | | | | | | | | | | | | | | |
| 20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS | | 21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED | | | | | | | | | | | | | | |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL Allen R. Miller | | 22b. TELEPHONE (<i>Include Area Code</i>) (202) 767-2215 | 22c. OFFICE SYMBOL Code 2330 | | | | | | | | | | | | | |

CONTENTS

| | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| INTRODUCTION | 1 |
| REDUCTION FORMULAE FOR $F_{q:1;0}^{p:2;1}$ $\left[\begin{matrix} \mu_1, \dots, \mu_p: \alpha, 1; 1; \\ \nu_1, \dots, \nu_q: \beta; -; x, x \end{matrix} \right]$ | 2 |
| REDUCTION FORMULAE FOR $F_{2:1;0}^{0:2;1}$ | 5 |
| PRELIMINARY RESULTS AND DEFINITIONS | 6 |
| ADDITIONAL RESULTS FOR $F_{2:1;0}^{0:2;1}$ | 8 |
| REFERENCES | 11 |

REDUCTION FORMULAE FOR KAMPE DE FERIET FUNCTIONS $F_{q:1;0}^{p:2;1}$

INTRODUCTION

In 1921, Kampé de Fériet defined and studied the functions named after him. Using the notation introduced in 1976 by Srivastava and Panda, the Kampé de Fériet functions in two variables are defined:

$$F_{l:m; n}^{p: q; k} \left[\begin{array}{c} (a_p); (b_q); (c_k); \\ (\alpha_l); (\beta_m); (\gamma_n); \end{array} x, y \right] := \sum_{r,s=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_{r+s} \prod_{j=1}^q (b_j)_r \prod_{j=1}^k (c_j)_s}{\prod_{j=1}^l (\alpha_j)_{r+s} \prod_{j=1}^m (\beta_j)_r \prod_{j=1}^n (\gamma_j)_s} \frac{x^r}{r!} \frac{y^s}{s!}.$$

The exact region of convergence for these functions is determined by using Horn's theorem for double series. Many properties and applications of Kampé de Fériet functions are found in Refs. 1-3.

There appears, however, to be a paucity of reduction formulae for Kampé de Fériet functions. Srivastava and Karlsson [1] list only 20 nontrivial instances in which Kampé de Fériet functions can be expressed in terms of generalized hypergeometric functions. In particular, for the functions $F_{q:1;0}^{p:2;1}$, they give the three results:

$$\begin{aligned} F_{q:1;0}^{p:2;1} & \left[\begin{array}{c} \alpha_1, \dots, \alpha_p; \lambda, \mu; v - \lambda - \mu; \\ \beta_1, \dots, \beta_q; v; \end{array} x, x \right] \\ & = {}_{p+2}F_{q+1} \left[\begin{array}{c} \alpha_1, \dots, \alpha_p, v - \lambda, v - \mu; \\ \beta_1, \dots, \beta_q, v; x \end{array} \right], \end{aligned} \quad (1)$$

$$\begin{aligned} F_{q:1;0}^{p:2;1} & \left[\begin{array}{c} \alpha_1, \dots, \alpha_p; \lambda, \mu; -1; \\ \beta_1, \dots, \beta_q; \lambda - \mu + 1; -; x, x \end{array} \right] \\ & = {}_{p+3}F_{q+2} \left[\begin{array}{c} \alpha_1, \dots, \alpha_p, \lambda - 1, \mu - 1, \frac{1}{2}\lambda + \frac{1}{2}; \\ \beta_1, \dots, \beta_q, \lambda - \mu + 1, \frac{1}{2}\lambda - \frac{1}{2}; x \end{array} \right], \end{aligned} \quad (2)$$

$$F_{q:0:0}^{p:1;1} \left[\begin{array}{c} \alpha_1, \dots, \alpha_p: \mu; \nu; \\ \beta_1, \dots, \beta_q: \frac{}{}; \frac{}{}; x, x \end{array} \right]$$

$$= {}_p F_q [\alpha_1, \dots, \alpha_p, \mu + \nu; \beta_1, \dots, \beta_q; x], \quad (3)$$

the latter being viewed as a special case of $F_{q:1:0}^{p:2;1}$.

In a series of recent papers [4-7], the author has shown that the function $F_{2:1:0}^{0:2;1}$ is intimately connected with representations for incomplete Lipschitz-Hankel integrals of cylindrical functions (see Eqs. (14-16)), where the class of cylindrical functions C includes Bessel functions of the first kind J , modified Bessel functions I , Bessel functions of the second kind or Neumann functions Y (or N), Bessel functions of imaginary argument or MacDonald functions K , and Bessel functions of the third kind that include Hankel functions of the first and second kind, $H^{(1)}$ and $H^{(2)}$.

As a byproduct of these investigations, several reduction formulae for $F_{2:1:0}^{0:2;1}$ not included in Eqs. (1) and (2) were derived. It is the purpose of this report, in addition to collecting these results, to generalize them somewhat and to derive four reduction formulae for $F_{2:1:0}^{0:2;1}$ that were not previously given.

REDUCTION FORMULAE FOR $F_{q:1:0}^{p:2;1}$ $\left[\begin{array}{c} \mu_1, \dots, \mu_p: \alpha, 1; 1; \\ \nu_1, \dots, \nu_q: \beta; -; \end{array} x, x \right]$

The following is derived in [4]:

$$\begin{aligned} & F_{2:1:0}^{0:2;1} \left[\begin{array}{c} \alpha, 1; 1; \\ \gamma, \delta: \beta; -; \end{array} x, x \right] \\ & = \frac{1 - \beta}{\alpha - \beta + 1} {}_1F_2[1; \gamma, \delta; x] + \frac{\alpha}{\alpha - \beta + 1} {}_2F_3[1, \alpha + 1; \gamma, \delta, \beta; x]. \end{aligned} \quad (4)$$

In particular, the following are special cases of Eq. (4) given essentially in terms of modified Bessel functions [5]:

$$\begin{aligned} & F_{2:1:0}^{0:2;1} \left[\begin{array}{c} 2 + \nu, 1; 1; \\ 3 + \nu, 3: 5/2; -; \end{array} \frac{x^2}{4}, \frac{x^2}{4} \right] \\ & = \frac{2 + \nu}{1 + 2\nu} \frac{48}{x^4} \left\{ \cosh x - 2 \Gamma(2 + \nu) \left(\frac{2}{x} \right)^{\nu} I_{\nu}(x) + 1 + 2\nu \right\}, \end{aligned} \quad (5)$$

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \text{---}: 1 + \nu, 1; -1; \frac{x^2}{4}, \frac{x^2}{4} \\ 2 + \nu, 2: \quad \quad \quad 3/2; -; \end{array} \right] = \frac{1 + \nu}{1 + 2\nu} \frac{4}{x^2} \left\{ \cosh x - \left(\frac{2}{x} \right)^\nu \Gamma(1 + \nu) I_\nu(x) \right\}, \quad (6)$$

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \text{---}: 1 + \nu, 1; -1; \frac{x^2}{4}, \frac{x^2}{4} \\ 1 + \nu, 2: \quad \quad \quad 3/2; -; \end{array} \right] = \frac{2}{1 + 2\nu} \frac{1}{x} \left\{ \frac{2\nu \cosh x}{x} + \sinh x - \left(\frac{2}{x} \right)^\nu \Gamma(1 + \nu) I_{\nu-1}(x) \right\}, \quad (7)$$

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \text{---}: 2 + \nu, 1; -1; \frac{x^2}{4}, \frac{x^2}{4} \\ 2 + \nu, 3: \quad \quad \quad 5/2; -; \end{array} \right] = \frac{1}{1 + 2\nu} \frac{24}{x^4} \left\{ x \sinh x + 2\nu \cosh x - 4\Gamma(2 + \nu) \left(\frac{2}{x} \right)^{\nu-1} I_{\nu-1}(x) + 2\nu(1 + 2\nu) \right\}. \quad (8)$$

Equation (4) is, in fact, included in the reduction formula

$$F_{q:1;0}^{p:2;1} \left[\begin{array}{c} \mu_1, \dots, \mu_p: \alpha, 1; -1; \\ \nu_1, \dots, \nu_q: \beta; -; x, x \end{array} \right] = \frac{1 - \beta}{1 - \beta + \alpha} {}_{p+1}F_q [\mu_1, \dots, \mu_p, 1; \nu_1, \dots, \nu_q; x] + \frac{\alpha}{1 - \beta + \alpha} {}_{p+2}F_{q+1} [\mu_1, \dots, \mu_p, 1, \alpha + 1; \nu_1, \dots, \nu_q, \beta; x], \quad (9)$$

which we now show.

By definition

$$\begin{aligned} F_{q:1;0}^{p:2;1} & \left[\begin{array}{c} \mu_1, \dots, \mu_p: \alpha, 1; -1; \\ \nu_1, \dots, \nu_q: \beta; -; x, x \end{array} \right] \\ &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\mu_1)_{m+n} \dots (\mu_p)_{m+n}}{(\nu_1)_{m+n} \dots (\nu_q)_{m+n}} \frac{(\alpha)_m (1)_m (1)_n}{(\beta)_m} \frac{x^m x^n}{m! n!} \\ &= \sum_{r=0}^{\infty} \left[\sum_{m=0}^r \frac{(\alpha)_m}{(\beta)_m} \right] \frac{(\mu_1)_r \dots (\mu_p)_r}{(\nu_1)_r \dots (\nu_q)_r} x^r. \end{aligned}$$

Since [8, Eq. (7.1.1), p. 151]

$$\sum_{m=0}^r \frac{(\alpha)_m}{(\beta)_m} = \frac{1}{1 - \beta + \alpha} \left[1 - \beta + \alpha \frac{(\alpha + 1)_r}{(\beta)_r} \right],$$

we arrive at

$$\begin{aligned} F_{q:1;0}^{p:2;1} & \left[\begin{array}{l} \mu_1, \dots, \mu_p : \alpha, 1; \quad 1; \\ \nu_1, \dots, \nu_q : \quad \beta; \quad -; \quad x, x \end{array} \right] \\ &= \frac{1 - \beta}{1 - \beta + \alpha} \sum_{r=0}^{\infty} \frac{(\mu_1)_r \dots (\mu_p)_r (1)_r}{(\nu_1)_r \dots (\nu_q)_r} \frac{x^r}{r!} \\ &+ \frac{\alpha}{1 - \beta + \alpha} \sum_{r=0}^{\infty} \frac{(\mu_1)_r \dots (\mu_p)_r}{(\nu_1)_r \dots (\nu_q)_r} \frac{(1)_r (\alpha + 1)_r}{(\beta)_r} \frac{x^r}{r!}, \end{aligned}$$

and Eq. (9) follows.

For brevity, we define below two functions that occur often in what follows:

$${}_p S_q [(a_p); (b_q); z]$$

$$:= \frac{1}{2} \{ e^z {}_p F_q [(a_p); (b_q); -2z] - e^{-z} {}_p F_q [(a_p); (b_q); 2z] \},$$

$${}_p C_q [(a_p); (b_q); z]$$

$$:= \frac{1}{2} \{ e^z {}_p F_q [(a_p); (b_q); -2z] + e^{-z} {}_p F_q [(a_p); (b_q); 2z] \}.$$

We observe that

$${}_p S_q [(a_p); (b_q); z] = \left\{ 1 - 2 \frac{\prod_{k=1}^p a_k}{\prod_{k=1}^q b_k} \right\} z + O(z^3) \quad z \rightarrow 0,$$

$${}_p C_q [(a_p); (b_q); z] = 1 + O(z^2) \quad z \rightarrow 0.$$

Hence,

$$\lim_{z \rightarrow 0} \frac{{}_p\mathbf{S}_q [(a_p); (b_q); z]}{z} = 1 - 2 \frac{\prod_{k=1}^p a_k}{\prod_{k=1}^q b_k},$$

$${}_p\mathbf{C}_q [(a_p); (b_q); 0] = 1.$$

REDUCTION FORMULAE FOR $F_{2:1;0}^{0:2;1}$

In addition, using the functions ${}_2\mathbf{C}_2$ and ${}_2\mathbf{S}_2$, we have from results in [6]

$$\begin{aligned} F_{2:1;0}^{0:2;1} &= {}_2\mathbf{C}_2 \left[\begin{array}{c} \frac{1 + \mu + \nu}{2}, \frac{2 + \mu + \nu}{2}; 1; \\ \frac{2 + \mu + \nu}{2}, \frac{3 + \mu + \nu}{2}; 1 + \nu; \end{array} \frac{x^2}{4}, \frac{x^2}{4} \right] \\ &= {}_2\mathbf{C}_2 \left[\begin{array}{c} 1 + \mu + \nu, 1/2 + \nu; \\ 2 + \mu + \nu, 1 + 2\nu; \end{array} x \right], \end{aligned} \quad (10)$$

$$\begin{aligned} F_{2:1;0}^{0:2;1} &= {}_2\mathbf{S}_2 \left[\begin{array}{c} \frac{1 + \mu + \nu}{2}, \frac{2 + \mu + \nu}{2}; 1; \\ \frac{3 + \mu + \nu}{2}, \frac{4 + \mu + \nu}{2}; 1 + \nu; \end{array} \frac{x^2}{4}, \frac{x^2}{4} \right] \\ &= \frac{2 + \mu + \nu}{2x} {}_2\mathbf{S}_2 \left[\begin{array}{c} 1 + \mu + \nu, 1/2 + \nu; \\ 2 + \mu + \nu, 1 + 2\nu; \end{array} x \right]. \end{aligned} \quad (11)$$

Equations (10) and (11) may also be written [6]

$$\begin{aligned}
 & F_{2:1;0}^{0:2;1} \left[\frac{1 + \mu + \nu}{2}, \frac{2 + \mu + \nu}{2}; 1; \frac{x^2}{4}, \frac{x^2}{4} \right] \\
 & = \cosh x {}_3F_4 \left[\frac{1 + \mu + \nu}{2}, \frac{1/2 + \nu}{2}, \frac{3/2 + \nu}{2}; \frac{x^2}{4} \right] \\
 & - \frac{1 + \mu + \nu}{2 + \mu + \nu} x \sinh x {}_3F_4 \left[\frac{2 + \mu + \nu}{2}, \frac{3/2 + \nu}{2}, \frac{5/2 + \nu}{2}; \frac{x^2}{4} \right], \quad (12)
 \end{aligned}$$

$$\begin{aligned}
 & F_{2:1;0}^{0:2;1} \left[\frac{1 + \mu + \nu}{2}, \frac{2 + \mu + \nu}{2}; 1; \frac{x^2}{4}, \frac{x^2}{4} \right] \\
 & = (2 + \mu + \nu) \frac{\sinh x}{x} {}_3F_4 \left[\frac{1 + \mu + \nu}{2}, \frac{1/2 + \nu}{2}, \frac{3/2 + \nu}{2}; \frac{x^2}{4} \right] \\
 & - (1 + \mu + \nu) \cosh x {}_3F_4 \left[\frac{2 + \mu + \nu}{2}, \frac{3/2 + \nu}{2}, \frac{5/2 + \nu}{2}; \frac{x^2}{4} \right] \quad (13)
 \end{aligned}$$

PRELIMINARY RESULTS AND DEFINITIONS

The general incomplete Lipschitz-Hankel integral of cylindrical functions $C_\nu(z)$ may be defined as the following function of two complex variables:

$$C_{e_{\mu,\nu}}(a, z) := \int_0^z e^{at} t^\mu C_\nu(t) dt. \quad (14)$$

Here the symbol e denotes the presence of the exponential function, and μ, ν may be complex. Analogously, we may define integrals that contain the functions $\sin(at)$ and $\cos(at)$ in place of $\exp(at)$:

$$C_{s_{\mu,\nu}}(a, z) := \int_0^z \sin(at) t^\mu C_\nu(t) dt, \quad (15)$$

$$C_{c_{\mu,\nu}}(a, z) := \int_0^z \cos(at) t^\mu C_\nu(t) dt. \quad (16)$$

To assure convergence of $C_{e_{\mu,\nu}}(a, z)$ and $C_{c_{\mu,\nu}}(a, z)$, it is necessary that $\operatorname{Re}(\mu + 1) > |\operatorname{Re} \nu|$ when $C = K, Y, H^{(1)}, H^{(2)}$; $\operatorname{Re}(1 + \mu + \nu) > 0$ when $C = I, J$. For convergence of $C_{s_{\mu,\nu}}(a, z)$, replace μ by $\mu + 1$ in the latter two inequalities.

Defining

$$\xi := \begin{cases} 1: C = I, K \\ -1: C = H, J, Y \end{cases} \quad \eta := \begin{cases} 1: C = K \\ -1: C = H, I, J, Y \end{cases}$$

$$Q[A_1(\mu, \nu); x, y] := F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 1}{2}; 1; \\ \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 3}{2}; 1/2; -; x, y \end{array} \right]$$

$$Q[A_2(\mu, \nu); x, y] := F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 1}{2}; 1; \\ \frac{\mu + \nu + 3}{2}, \frac{\mu - \nu + 3}{2}; 1/2; -; x, y \end{array} \right]$$

$$Q[B_1(\mu, \nu); x, y] := F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 2}{2}; 1; \\ \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 4}{2}; 3/2; -; x, y \end{array} \right]$$

$$Q[B_2(\mu, \nu); x, y] := F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 2}{2}; 1; \\ \frac{\mu + \nu + 4}{2}, \frac{\mu - \nu + 4}{2}; 3/2; -; x, y \end{array} \right],$$

it is shown in [7] that

$$C_{s_{\mu,\nu}}(a, z) = \frac{az^{2+\mu}}{\mu-\nu+2} \left\{ C_\nu(z) Q \left[B_1; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] + \frac{\eta z}{\mu+\nu+2} C_{\nu-1}(z) Q \left[B_2; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\}, \quad (17)$$

$$C_{c_{\mu,\nu}}(a, z) = \frac{z^{1+\mu}}{\mu-\nu+1} \left\{ C_\nu(z) Q \left[A_1; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] + \frac{\eta z}{\mu+\nu+1} C_{\nu-1}(z) Q \left[A_2; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\}. \quad (18)$$

ADDITIONAL RESULTS FOR $F_{2:1;0}^{0:2;1}$

In Eq. (17), set firstly $\xi = 1$, $\eta = -1$, $a = i$ and secondly $\xi = 1$, $\eta = 1$, $a = i$. This gives the system of linear equations in two unknowns $Q[B_1; z^2/4, z^2/4]$ and $Q[B_2; z^2/4, z^2/4]$:

$$\begin{aligned} I_\nu(z) Q \left[B_1; \frac{z^2}{4}, \frac{z^2}{4} \right] - \frac{z}{\mu + \nu + 2} I_{\nu-1}(z) Q \left[B_2; \frac{z^2}{4}, \frac{z^2}{4} \right] \\ = \frac{\mu - \nu + 2}{z^{2+\mu}} \int_0^z \sinh t \ t^\mu I_\nu(t) dt, \end{aligned} \quad (19)$$

$$\begin{aligned} K_\nu(z) Q \left[B_1; \frac{z^2}{4}, \frac{z^2}{4} \right] + \frac{z}{\mu + \nu + 2} K_{\nu-1}(z) Q \left[B_2; \frac{z^2}{4}, \frac{z^2}{4} \right] \\ = \frac{\mu - \nu + 2}{z^{2+\mu}} \int_0^z \sinh t \ t^\mu K_\nu(t) dt. \end{aligned} \quad (20)$$

Now making the previous substitutions in Eq. (18) gives the system of linear equations in unknowns $Q[A_1; z^2/4, z^2/4]$ and $Q[A_2; z^2/4, z^2/4]$:

$$\begin{aligned} I_\nu(z) Q \left[A_1; \frac{z^2}{4}, \frac{z^2}{4} \right] - \frac{z}{\mu + \nu + 1} I_{\nu-1}(z) Q \left[A_2; \frac{z^2}{4}, \frac{z^2}{4} \right] \\ = \frac{\mu - \nu + 1}{z^{1+\mu}} \int_0^z \cosh t \ t^\mu I_\nu(t) dt, \end{aligned} \quad (21)$$

$$\begin{aligned} K_\nu(z) Q \left[A_1; \frac{z^2}{4}, \frac{z^2}{4} \right] + \frac{z}{\mu + \nu + 1} K_{\nu-1}(z) Q \left[A_2; \frac{z^2}{4}, \frac{z^2}{4} \right] \\ = \frac{\mu - \nu + 1}{z^{1+\mu}} \int_0^z \cosh t \ t^\mu K_\nu(t) dt. \end{aligned} \quad (22)$$

Solving the systems Eqs. (19) and (20), and Eqs. (21) and (22), respectively, using Cramer's rule and noting the Wronskian

$$I_\nu(z) K_{\nu-1}(z) + I_{\nu-1}(z) K_\nu(z) = 1/z, \quad (23)$$

we obtain

$$Q \left[B_1; \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{\mu - \nu + 2}{z^{1+\mu}} \begin{vmatrix} \int_0^z \sinh t \ t^\mu I_\nu(t) dt & - I_{\nu-1}(z) \\ \int_0^z \sinh t \ t^\mu K_\nu(t) dt & K_{\nu-1}(z) \end{vmatrix}, \quad (24)$$

$$Q\left[B_2; \frac{z^2}{4}, \frac{z^2}{4}\right] = \frac{(\mu - \nu + 2)(\mu + \nu + 2)}{z^{2+\mu}} \begin{vmatrix} I_\nu(z) & \int_0^z \sinh t t^\mu I_\nu(t) dt \\ K_\nu(z) & \int_0^z \sinh t t^\mu K_\nu(t) dt \end{vmatrix}, \quad (25)$$

$$Q\left[A_1; \frac{z^2}{4}, \frac{z^2}{4}\right] = \frac{(\mu - \nu + 1)}{z^\mu} \begin{vmatrix} \int_0^z \cosh t t^\mu I_\nu(t) dt & -I_{\nu-1}(z) \\ \int_0^z \cosh t t^\mu K_\nu(t) dt & K_{\nu-1}(z) \end{vmatrix}, \quad (26)$$

$$Q\left[A_2; \frac{z^2}{4}, \frac{z^2}{4}\right] = \frac{(\mu - \nu + 1)(\mu + \nu + 1)}{z^{1+\mu}} \begin{vmatrix} I_\nu(z) & \int_0^z \cosh t t^\mu I_\nu(t) dt \\ K_\nu(z) & \int_0^z \cosh t t^\mu K_\nu(t) dt \end{vmatrix}. \quad (27)$$

Equations (24) and (25) are valid for $\operatorname{Re}(\mu \pm \nu + 2) > 0$; Eqs. (26) and (27) are valid for $\operatorname{Re}(\mu \pm \nu + 1) > 0$. Further, Eqs. (24-27) provide integral representations for the function Q appearing in each equation.

Using [9, p. 117, Eq. (2)], we evaluate the four integrals in Eqs. (24-27) in terms of the functions ${}_2C_2$ and ${}_2S_2$:

$$\begin{aligned} \int_0^z \sinh t t^\mu C_\nu(t) dt &= \frac{z^{\mu+1}}{\mu + \nu + 1} \left\{ \sinh z C_\nu(z) \right. \\ &\quad \left. - \frac{z C_\nu(z)}{\mu + \nu + 2} {}_2C_2 \left[\begin{matrix} 1, & \mu + 3/2; \\ \mu - \nu + 1, & \mu + \nu + 3; \end{matrix} z \right] + \frac{\delta z C_{\nu+1}(z)}{\mu - \nu + 1} {}_2S_2 \left[\begin{matrix} 1, & \mu + 3/2; \\ \mu - \nu + 2, & \mu + \nu + 2; \end{matrix} z \right] \right\}, \\ \int_0^z \cosh t t^\mu C_\nu(t) dt &= \frac{z^{1+\mu}}{\mu + \nu + 1} \left\{ \cosh z C_\nu(z) \right. \\ &\quad \left. - \frac{z C_\nu(z)}{\mu + \nu + 2} {}_2S_2 \left[\begin{matrix} 1, & \mu + 3/2; \\ \mu - \nu + 1, & \mu + \nu + 3; \end{matrix} z \right] + \frac{\delta z C_{\nu+1}(z)}{\mu - \nu + 1} {}_2C_2 \left[\begin{matrix} 1, & \mu + 3/2; \\ \mu - \nu + 2, & \mu + \nu + 2; \end{matrix} z \right] \right\}, \end{aligned}$$

where $\delta = 1$ if $C = K$, $\delta = -1$ if $C = I$.

Using these results in Eqs. (24-27), the F -symbol for Q and taking note of Eq. (23) together with the easily proved identity

$$K_{\nu+1}(z) I_{\nu-1}(z) - I_{\nu+1}(z) K_{\nu-1}(z) = 2\nu/z^2,$$

we deduce the four reduction formulae:

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 2}{2}; 1; \\ \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 4}{2} : \\ \end{array} \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{\mu - \nu + 2}{\mu + \nu + 1} \left\{ \frac{\sinh z}{z} - \frac{1}{\mu + \nu + 2} {}_2C_2 \left[\begin{array}{c} 1, \mu + 3/2; \\ \mu - \nu + 1, \mu + \nu + 3; \end{array} z \right] + \frac{2\nu}{\mu - \nu + 1} \frac{1}{z} {}_2S_2 \left[\begin{array}{c} 1, \mu + 3/2; \\ \mu - \nu + 2, \mu + \nu + 2; \end{array} z \right] \right\}, \quad (28)$$

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 2}{2}; 1; \\ \frac{\mu + \nu + 4}{2}, \frac{\mu - \nu + 4}{2} : \\ \end{array} \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{(\mu + \nu + 2)(\mu - \nu + 2)}{(\mu + \nu + 1)(\mu - \nu + 1)} \frac{1}{z} {}_2S_2 \left[\begin{array}{c} 1, \mu + 3/2; \\ \mu - \nu + 2, \mu + \nu + 2; \end{array} z \right], \quad (29)$$

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 1}{2}; 1; \\ \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 3}{2} : \\ \end{array} \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{\mu - \nu + 1}{\mu + \nu + 1} \left\{ \cosh z - \frac{z}{\mu + \nu + 2} {}_2S_2 \left[\begin{array}{c} 1, \mu + 3/2; \\ \mu - \nu + 1, \mu + \nu + 3; \end{array} z \right] + \frac{2\nu}{\mu - \nu + 1} {}_2C_2 \left[\begin{array}{c} 1, \mu + 3/2; \\ \mu - \nu + 2, \mu + \nu + 2; \end{array} z \right] \right\}, \quad (30)$$

$$F_{2:1;0}^{0:2;1} \left[\begin{array}{c} \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 1}{2}; 1; \\ \frac{\mu + \nu + 3}{2}, \frac{\mu - \nu + 3}{2} : \\ \end{array} \frac{z^2}{4}, \frac{z^2}{4} \right] = {}_2C_2 \left[\begin{array}{c} 1, \mu + 3/2; \\ \mu - \nu + 2, \mu + \nu + 2; \end{array} z \right]. \quad (31)$$

If we set $\mu = \nu$, then it is easily shown that Eq. (28) reduces to Eq. (7), and Eq. (29) reduces to Eq. (6). Equations (30) and (31) reduce to $F_{2:0;0}^{0:1;1}$ functions (see Eqs. (22) and (23) of Ref. 5), whose reduction is given by Eq. (3).

We remark that Eqs. (28-31) are valid for $|z| < \infty$ and for all μ, ν such that each function exists. Hence the restrictions on the existence of the integrals in Eqs. (24-27) do not apply here.

REFERENCES

1. H.M. Srivastava and P.W. Karlsson, *Multiple Gaussian Hypergeometric Series* (Ellis Horwood Ltd., Chichester, U.K., 1985).
2. H.M. Srivastava and H.L. Manocha, *A Treatise on Generating Functions* (Ellis Horwood Ltd., Chichester, U.K., 1984).
3. H. Exton, *Handbook of Hypergeometric Integrals* (Ellis Horwood Ltd., Chichester, U.K., 1978).
4. A.R. Miller, "A Lipschitz-Hankel Integral of K_0 ," *J. Franklin Institute*, **325**, No. 6, pp. 787-795, 1988.
5. A.R. Miller, "Incomplete Lipschitz-Hankel Integrals of MacDonald Functions," *J. Math. Anal. Appl.* (to appear). (Or see NRL Report 9112, March 1988.)
6. A.R. Miller, "Incomplete Lipschitz-Hankel Integrals of Bessel Functions," *J. Math. Anal. Appl.* (to appear). (Or see NRL Report 9047, May 1987.)
7. A.R. Miller, "A Note on Incomplete Integrals of Cylindrical Functions," *J. Franklin Institute* (to appear). (Or see NRL Report 9127, June 1988.)
8. E.R. Hansen, *A Table of Series and Products* (Prentice-Hall, Englewood Cliffs, N.J., 1975).
9. Y.L. Luke, *Integrals of Bessel Functions* (McGraw-Hill, New York, 1962).

